

Abelian functions satisfy an Algebraic Addition Theorem

Mark B. Villarino

Escuela de Matemática, Universidad de Costa Rica,
2060 San José, Costa Rica

April 16, 1998

Abstract

We prove that n independent abelian functions admit an algebraic addition theorem, with no appeal to theta functions.

A famous and classical theorem of Weierstrass states:

Theorem 1 (Weierstrass, 1869). *Any n analytically independent Abelian functions in n complex variables that belong to the same $2n$ -dimensional period lattice satisfy an algebraic addition theorem.*

All published proofs of this theorem depend on the Weierstrass–Riemann “Theta-satz” which affirms that any abelian function is globally representable as the quotient of two (generalized) theta functions [2].

We show in this note that the theorem has nothing to do with theta functions *per se*, but is a simple (indeed, almost trivial) consequence of another, even more famous, theorem of Weierstrass [4], Thimm [3] and Siegel [1] on algebraic dependence.

Theorem 2 (Weierstrass–Thimm–Siegel). *Any $n + 1$ meromorphic functions on a compact complex space of dimension n are algebraically dependent.*

Definition 1. A set of n meromorphic functions of n complex variables, $\varphi_k(u_1, \dots, u_n)$, $k = 1, \dots, n$, satisfy an **algebraic addition theorem** if and only if there exist n polynomials $G_1, \dots, G_n$ in $2n + 1$ complex variables with complex coefficients, such that the equations

$$G_k[\varphi_k(\mathbf{u} + \mathbf{v}); \varphi_1(\mathbf{u}), \dots, \varphi_n(\mathbf{u}); \varphi_1(\mathbf{v}), \dots, \varphi_n(\mathbf{v})] = 0$$

for $k = 1, \dots, n$, hold identically for all $\mathbf{u} = (u_1, \dots, u_n) \in \mathbb{C}^n$ and $\mathbf{v} = (v_1, \dots, v_n) \in \mathbb{C}^n$ where the φ_i are defined.

Definition 2. The meromorphic functions $\varphi_1, \dots, \varphi_k$ on the compact complex space X are called **algebraically dependent** iff there exists a polynomial $Q(z_1, \dots, z_k) \not\equiv 0$ with complex coefficients such that

$$Q(\varphi_1(\mathbf{u}), \dots, \varphi_k(\mathbf{u})) = 0$$

holds identically for all $\mathbf{u} \in X$ where the φ_k are defined.

Proof of Theorem 1. Let X be the period parallelootope for the lattice Ω of periods of the abelian functions $\varphi_1(\mathbf{u}), \dots, \varphi_n(\mathbf{u})$. Then the cartesian product $X \times X$ is a compact complex connected manifold of dimension $2n$.

The function $\varphi_k(\mathbf{u} + \mathbf{v})$ is a meromorphic function of $2n$ variables $(u_1, \dots, u_n, v_1, \dots, v_n)$ on $X \times X$, as are the $2n$ functions $\varphi_k(\mathbf{u})$ and $\varphi_k(\mathbf{v})$. Therefore, the $2n + 1$ meromorphic functions $\varphi_k(\mathbf{u} + \mathbf{v})$; $\varphi_1(\mathbf{u}), \dots, \varphi_n(\mathbf{u})$; $\varphi_1(\mathbf{v}), \dots, \varphi_n(\mathbf{v})$ on the $2n$ -dimensional compact complex space $X \times X$ are necessarily algebraically dependent, by Theorem 2. This conclusion holds for each $\varphi_k(\mathbf{u} + \mathbf{v})$, $k = 1, \dots, n$. But, by Definition 1, this means that $\varphi_1, \dots, \varphi_k$ satisfy an algebraic addition theorem. $\square$

References

- [1] C. L. Siegel, “Meromorphe Funktionen auf kompakten analytischen Mannigfaltigkeiten”, *Nachr. Akad. Wiss. Göttingen* **4** (1955), 71–77.
- [2] C. L. Siegel, *Lectures on Complex Function Theory III*, Addison-Wesley, Reading, MA, 1972.
- [3] W. Thimm, “Meromorphe Abbildungen von Riemannschen Bereichen”, *Math. Z.* **60** (1954), 435–457.
- [4] K. Weierstrass, “Untersuchungen über die $2r$ -fach periodischen Funktionen von r Veränderlichen”, *J. reine angew. Math.* **89** (1880), 1–8. (Collected Works, vol. II, pp. 125–133.)